



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

Mathematical Social Sciences 48 (2004) 11–22

mathematical  
social  
sciences

[www.elsevier.com/locate/econbase](http://www.elsevier.com/locate/econbase)

# A note on Wakker's Cardinal Coordinate Independence

Denis Bouyssou<sup>a,\*</sup>, Marc Pirlot<sup>b</sup>

<sup>a</sup> *CNRS, LAMSADE, Université Paris Dauphine, Place du Maréchal de Lattre de Tassigny, F-75775, Paris Cedex 16, France*

<sup>b</sup> *Faculté Polytechnique de Mons, 9, rue de Houdain, B-7000 Mons, Belgium*

Received 30 August 2003; received in revised form 1 December 2003; accepted 13 January 2004

---

## Abstract

Peter P. Wakker has forcefully shown the importance for decision theory of a condition that he called “Cardinal Coordinate Independence” (CCI). Indeed, when the outcome space is rich, he proved that, for continuous weak orders, this condition fully characterizes the Subjective Expected Utility (SEU) model with a finite number of states. He has furthermore explored in depth how this condition can be weakened in order to arrive at characterizations of Choquet Expected Utility and Cumulative Prospect Theory. This note studies the consequences of this condition in the absence of any transitivity assumption. Complete preference relations satisfying Cardinal Coordinate Independence are shown to be already rather well-behaved. Under a suitable necessary order denseness assumption, they may always be represented using a simple numerical model.

© 2004 Elsevier B.V. All rights reserved.

*Keywords:* Decision under uncertainty; Cardinal Coordinate Independence; Nontransitive preferences

*JEL classification:* D81

---

## 1. Introduction and motivation

The work of Peter P. Wakker on the foundations of decision theory has forcefully shown how the consideration of induced relations comparing preference differences

---

\*Corresponding author. Tel.: +33-1-44054898; fax: +33-1-44054091.

*E-mail addresses:* [bouyssou@lamsade.dauphine.fr](mailto:bouyssou@lamsade.dauphine.fr) (D. Bouyssou), [marc.pirlot@fpms.ac.be](mailto:marc.pirlot@fpms.ac.be) (M. Pirlot).

<sup>1</sup>Part of this work was accomplished while Denis Bouyssou was visiting the Service de Mathématique de la Gestion at the Université Libre de Bruxelles (Brussels, Belgium). The warm hospitality of the Service de Mathématique de la Gestion, the support of the Belgian Fonds National de la Recherche Scientifique and the Brussels-Capital Region through a “Research in Brussels” action grant are gratefully acknowledged.

between outcomes may illuminate the analysis of models of decision making under uncertainty.<sup>2</sup> In order to characterize the Subjective Expected Utility (SEU) model with a finite number of states, he introduced a condition called “Cardinal Coordinate Independence” (CCI)<sup>3</sup>. This condition requires that the comparisons of preference differences between outcomes revealed in different states and using different reference outcomes do not exhibit contradictory information. Using topological assumptions to ensure that the set of outcomes is “rich”, Wakker showed that CCI fully characterizes SEU for continuous weak orders (see Wakker, 1984, 1988b, 1989a). This striking result can be extended to more general outcome sets in the algebraic approach (see Wakker, 1991).

This condition, when appropriately weakened (e.g. requiring it only for comonotonic acts), may also be used to characterize non-EU models such as Choquet Expected Utility (see Wakker, 1989a,b, 1994) and Cumulative Prospect Theory (Wakker and Tversky, 1993). Indeed, CCI and its variants may be seen as a powerful unifying tool to analyze many models in decision making under risk and uncertainty (see Wakker and Zank, 1999). Furthermore, this condition is intimately related to empirical assessment techniques of utility functions and has served as an inspiring principle for the development of such techniques (see Abdellaoui, 2000; Bleichrodt and Pinto, 2000; Wakker and Deneffe, 1996).

In all the above-mentioned papers, it is supposed that the set of outcomes is rich and that the complete and transitive preferences behave consistently in this rich structure. This is done either requiring a “solvability” assumption (see Wakker, 1988a, 1991) or continuity w.r.t. a connected topology (see Wakker, 1988a, 1991). Clearly, these structural assumptions interact with the necessary conditions and it is well-known that this may contribute to obscure their interpretation (see Furkhen and Richter, 1991; Krantz et al., 1971, ch. 9, Wakker, 1989a, pages 75–76, or Köbberling and Wakker, 2003, Appendix A). Furthermore, the weak order assumption is quite powerful: transitivity clearly plays a vital role in the definition of “standard sequences” or “grids” (see, e.g. Krantz et al., 1971; Wakker, 1989a). In view of the importance of CCI, it seems worth investigating its “pure consequences”, i.e. its consequences in the absence of any transitivity requirement and of any unnecessary structural assumption on the set of outcomes. This is the purpose of this note.

Rather surprisingly, it turns out that, when coupled with completeness, CCI already implies that preferences are rather well-behaved. Under a suitable necessary order denseness assumption, such preferences may be represented numerically using a simple model that generalizes the Skew Symmetric Additive (SSA) model studied by Fishburn (1990), replacing additivity by a mere decomposability requirement. Technically (so to speak, since in our poor framework, the reader should not expect anything that is much involved), our results extend to the case of decision making under uncertainty the results obtained by Bouyssou and Pirlot (2002) in the case of conjoint measurement.

In this note, models tolerating intransitive preferences are simply used as a framework allowing to understand the pure consequences of some well-known conditions. They may nevertheless have some interest in themselves. Indeed, as shown by the famous experiment

<sup>2</sup>A similar idea is already used by Pfanzagl (1971, ch. 9); we thank Peter Wakker for bringing this point to our attention.

<sup>3</sup>We use here the terminology of Wakker (1984). This condition is often called “tradeoff consistency” in Wakker’s later texts, e.g. Wakker (1988b, 1989a).

in Tversky (1969), nontransitive preferences may be observed in quite a predictable way in decision making under risk (see, however, Iveson and Falmagne (1985), for a critical analysis of this experiment). Furthermore, Fishburn (1991) has challenged, quite convincingly in our opinion, the usual arguments used to dismiss such models (for classical and less classical counterarguments, see Luce, 2000; Raiffa, 1968).

This note is organized as follows. Section 2 briefly introduces our setting and notation. Our main results appear in Section 3. A final section discusses our findings.

## 2. The setting

### 2.1. Outcomes, states and acts

We adopt a classical setting for decision under uncertainty with a finite number of states. Let  $\Gamma = \{\alpha, \beta, \gamma, \dots\}$  be the set of *outcomes* and  $N = \{1, 2, \dots, n\}$  be the set of *states*. It is understood that the elements of  $N$  are exhaustive and mutually exclusive: one and only one state will turn out to be true. An *act* is a function from  $N$  to  $\Gamma$ . The set of all acts is denoted by  $\mathcal{A} = \Gamma^N$ . Acts will always be denoted by lowercase letters  $a, b, c, d, \dots$ . An act  $a \in \mathcal{A}$  therefore associates to each state  $i \in N$  an outcome  $a(i) \in \Gamma$ . We often abuse notation and write  $a_i$  instead of  $a(i)$ . Among the elements of  $\mathcal{A}$  are constant acts, i.e. acts giving the same outcome in all states. We denote by  $\bar{\alpha}$  the constant act giving the outcome  $\alpha \in \Gamma$  in all states  $i \in N$ .

Let  $E \subseteq N$  and  $a, b \in \mathcal{A}$ . We denote by  $a_E b$  the act  $c \in \mathcal{A}$  such that  $c_i = a_i$ , for all  $i \in E$  and  $c_i = b_i$ , for all  $i \in N \setminus E$ . Similarly  $\alpha_E b$  will denote the act  $d \in \mathcal{A}$  such that  $d_i = \alpha$ , for all  $i \in E$  and  $d_i = b_i$ , for all  $i \in N \setminus E$ . When  $E = \{i\}$ , we write  $a_i b$  and  $\alpha_i b$  instead of  $a_{\{i\}} b$  and  $\alpha_{\{i\}} b$ .

### 2.2. Preferences on acts

In this note,  $\succeq$  will always denote a reflexive binary relation on the set  $\mathcal{A}$ . The binary relation  $\succeq$  is interpreted as an “at least as good as” preference relation between acts. We denote by  $\succ$  (resp.  $\sim$ ) the asymmetric (resp. symmetric) part of  $\succeq$ . A similar convention holds when  $\succeq$  is starred, superscripted and/or subscripted.

We say that state  $i \in N$  is *influential* (for  $\succeq$ ) if there are  $\alpha, \beta, \gamma, \delta \in \Gamma$  and  $a, b \in \mathcal{A}$  such that  $\alpha_i a \succeq \beta_i b$  and  $\text{Not}[\gamma_i a \succeq \delta_i b]$  and degenerate otherwise. It is clear that a degenerate state has no influence whatsoever on the comparison of the elements of  $\mathcal{A}$  and may be suppressed from  $N$ . In order to avoid unnecessary minor complications, we suppose henceforth that all states in  $N$  are influential. Note that, in general, this does not rule out the existence of null states  $i \in N$ , i.e. such that  $a_i c \sim b_i c$ , for all  $a, b, c \in \mathcal{A}$ . A state will be said *essential* if it is not null.<sup>4</sup>

We denote by  $\succeq_\Gamma$  the relation induced on the set of outcomes  $\Gamma$  by the relation  $\succeq$  on  $\mathcal{A}$ , i.e., for all  $\alpha, \beta \in \Gamma$ ,  $\alpha \succeq_\Gamma \beta \Leftrightarrow \bar{\alpha} \succeq \bar{\beta}$ .

<sup>4</sup>As pointed out to us by a referee, the distinction between an influential and an essential state is subtle. Let us illustrate this difference with the simple example of a preference relation for which each state is null while all states are influential. Let  $N = \{1, 2, 3, 4\}$  and  $\Gamma = \mathbb{R}$ . Let  $\succeq$  on  $\mathcal{A}$  be such that, for all  $a, b \in \mathcal{A}$ ,  $a \succeq b \Leftrightarrow \sum_{i \in P(a,b)} p_i \geq \sum_{i \in P(b,a)} p_i - 1/4$ , with  $p_1 = p_2 = p_3 = p_4 = 1/4$  and  $P(a,b) = \{i \in N: a_i > b_i\}$ . The relation  $\succeq$  is clearly complete. All states are null (an essential state would require to have  $p_i > 1/4$ ). Yet, each state is influential. For instance, state 1 is influential because  $\text{Not}[(0, 0, 0, 0) \succeq (10, 10, 0, 0)]$  but  $(10, 0, 0, 0) \succeq (10, 10, 0, 0)$ .

### 2.3. Comparing preference differences between outcomes

Consistently with Wakker (1988b, 1989a), our analysis uses induced relations comparing preference differences on the set of outcomes. Note that our definitions differ from his, although we use similar notation.

The idea that any binary relation generates various reflexive and transitive binary relations called traces dates back at least to the pioneering work of Luce (1956). The use of traces has proved especially useful in the study of preference structures tolerating intransitive indifference such as semiorders or interval orders (see Aleskerov and Monjardet, 2002; Fishburn, 1985; Pirlot and Vincke, 1997). We pursue here the same idea using traces on preference differences.

**Definition 1.** (Relations comparing preference differences). Let  $\succeq$  be a binary relation on  $\mathcal{A}$ . We define the binary relations  $\succeq^*$  and  $\succeq^{**}$  on  $\Gamma^2$  letting, for all  $\alpha, \beta, \gamma, \delta \in \Gamma$ ,

$$\begin{aligned}
 (\alpha, \beta) \succeq^*(\gamma, \delta) & \text{ if [for all } a, b \in \mathcal{A} \text{ and all } i \in N, \gamma_i a \succeq \delta_i b \Rightarrow \alpha_i a \succeq \beta_i b], \\
 (\alpha, \beta) \succeq^{**}(\gamma, \delta) & \text{ if } [(\alpha, \beta) \succeq^*(\gamma, \delta) \text{ and } (\delta, \gamma) \succeq^*(\beta, \alpha)].
 \end{aligned}$$

Intuitively, if  $(\alpha, \beta) \succeq^*(\gamma, \delta)$ , it seems reasonable to conclude that the preference difference between  $\alpha$  and  $\beta$  is not smaller than the preference difference between  $\gamma$  and  $\delta$ . Contrary to the intuition concerning the comparison of preference differences, the definition of  $\succeq^*$  does not imply that the two “opposite” differences  $(\alpha, \beta)$  and  $(\beta, \alpha)$  are linked. This is the motivation for the introduction of the relation  $\succeq^{**}$ .

By construction,  $\succeq^*$  and  $\succeq^{**}$  are reflexive and transitive. Therefore,  $\sim^*$  and  $\sim^{**}$  are equivalence relations. Note that, by construction,  $\succeq^{**}$  is reversible, i.e.  $(\alpha, \beta) \succeq^{**}(\gamma, \delta) \Leftrightarrow (\delta, \gamma) \succeq^{**}(\beta, \alpha)$ . Observe that  $\succeq^*$  and  $\succeq^{**}$  may not be complete: induced comparisons of preference differences may indeed depend on the reference acts and/or on the state in which they are revealed. Sweeping consequences will obtain when this is not so.

We note a few useful connections between  $\succeq^*$ ,  $\succeq^{**}$  and  $\succeq$  in the following lemma that holds independently on any condition on  $\succeq$ .

**Lemma 2.** *Let  $\succeq$  be a binary relation on  $\mathcal{A}$ . Then, for all  $a, b, c, d \in \mathcal{A}$  and all  $i \in N$ ,*

$$[a \succeq b \text{ and } (c_i, d_i) \succeq^*(a_i, b_i)] \Rightarrow c_i a \succeq d_i b \tag{1}$$

$$[(c_j, d_j) \sim^*(a_j, b_j), \text{ for all } j \in N] \Rightarrow [a \succeq b \Leftrightarrow c \succeq d], \tag{2}$$

$$[a \succ b \text{ and } (c_i, d_i) \succeq^{**}(a_i, b_i)] \Rightarrow c_i a \succ d_i b, \tag{3}$$

$$[(c_j, d_j) \sim^{**}(a_j, b_j), \text{ for all } j \in N] \Rightarrow \begin{cases} a \succeq b \Leftrightarrow c \succeq d \text{ and} \\ a \succ b \Leftrightarrow c \succ d. \end{cases} \tag{4}$$

**Proof.** Eq. (1) is obvious from the definition of  $\succeq^*$ , Eq. (2) follows. Suppose that  $a \succ b$  and  $(c_i, d_i) \succeq^{**}(a_i, b_i)$ . Using Eq. (1), we know that  $c_i a \succeq d_i b$ . Suppose now that  $d_i b \succeq c_i a$ . Since  $(c_i, d_i) \succeq^{**}(a_i, b_i)$  implies  $(b_i, a_i) \succeq^*(d_i, c_i)$ , Eq. (1) implies  $b \succeq a$ , a contradiction. Hence Eq. (3) holds and Eq. (4) follows.  $\square$

### 2.4. Coordinate independence and Cardinal Coordinate Independence

Coordinate Independence (CI) is a classical independence condition stating that the preference between acts is not affected by a common outcome in some state.

**Definition 3.** (Condition CI). Let  $\succeq$  be a binary relation on  $\mathcal{A}$ . This relation is said to satisfy CI if, for all  $i \in N$ , all  $\alpha, \beta \in \Gamma$  and all  $a, b \in \mathcal{A}$ ,

$$\alpha_i a \succeq \alpha_i b \Rightarrow \beta_i a \succeq \beta_i b.$$

It is not difficult to see that if  $\succeq$  satisfies CI then, for all  $E \subset N$  and all  $a, b, c, d \in \mathcal{A}$ ,  $a_E c \succeq b_E c \Rightarrow a_E d \succeq b_E d$ . Condition CI is therefore equivalent to postulate P2, often referred to as the “Sure Thing Principle”, introduced by [Savage \(1954\)](#).

The following definition is adapted from [Wakker \(1989a, page 80\)](#).

**Definition 4.** (Condition CCI). Let  $\succeq$  be a binary relation on  $\mathcal{A}$ . This relation is said to satisfy CCI if:

$$\left. \begin{array}{l} \alpha_i a \succeq \beta_i b \\ \text{and} \\ \gamma_i b \succeq \delta_i a \\ \text{and} \\ \delta_j c \succeq \gamma_j d \end{array} \right\} \Rightarrow \alpha_j c \succeq \beta_j d,$$

for all  $i, j \in N$ , all  $a, b, c, d \in \mathcal{A}$  and all  $\alpha, \beta, \gamma, \delta \in \Gamma$ .

We refer to [Wakker \(1989a\)](#) for a thorough discussion of this condition and to the references in Section 1 for a study of its possible variants.<sup>5</sup> Note that, since we supposed that all degenerate states were suppressed from  $N$ , we state here the condition for all  $i, j \in N$ , contrary to [Wakker \(1989a\)](#) who only states it for essential states  $i$  and  $j$ . Although influent states may not be essential, they will turn out to be so for complete relations in presence of CCI.

## 3. Results

### 3.1. CI, CCI and preference differences

Conditions CI and CCI can easily be reformulated using the relations comparing preference differences between outcomes introduced above. The, easy, case of CI is dealt with first.

<sup>5</sup>[Köbberling and Wakker \(2003\)](#) have recently proposed a weakened version of CCI, using indifferences instead of preferences in the above definition. They show that this condition, when coupled with monotonicity, may replace the original condition in the characterization of SEU (they furthermore show that such a weakening is also possible starting with the restricted versions of CCI mentioned above used to characterize Choquet Expected Utility and Cumulative Prospect Theory). In our nontransitive setting, such a weakening of CCI does not seem to lead, on its own, to interesting restrictions on  $\succeq$ . We do not consider it here.

**Proposition 5.** Let  $\succeq$  be a binary relation on  $\mathcal{A}$ . Then  $\succeq$  satisfies CI if and only if and only if (iff)  $(\alpha, \alpha) \sim^*(\beta, \beta)$ , for all  $\alpha, \beta \in \Gamma$ .

**Proof.** It is clear that  $[\succeq \text{ satisfies CI}] \Leftrightarrow [\alpha_i a \succeq \alpha_i b \Leftrightarrow \beta_i a \succeq \beta_i b, \text{ for all } \alpha, \beta \in \Gamma, \text{ all } i \in N \text{ and all } a, b \in \mathcal{A}] \Leftrightarrow [(\alpha, \alpha) \sim^*(\beta, \beta) \text{ for all } \alpha, \beta \in \Gamma]. \quad \square$

The following summarizes the main consequences of CCI when  $\succeq$  is complete.

**Lemma 6.** Let  $\succeq$  be a complete relation on  $\mathcal{A}$ . If  $\succeq$  satisfies CCI then:

- (1)  $\succeq^*$  is complete,
- (2)  $\text{Not}[(\alpha, \beta) \succeq^*(\gamma, \delta)] \Rightarrow (\beta, \alpha) \succeq^*(\delta, \gamma)$ ,
- (3)  $\succeq^{**}$  is complete,
- (4)  $\succeq$  satisfies CI,
- (5)  $[a \sim b \text{ and } (\gamma, \delta) \succ^{**}(a_i, b_i)] \Rightarrow \gamma_i a \succ \delta_i b$ ,
- (6) all states are essential.

for all  $a, b \in \mathcal{A}$ , all  $i \in N$  and all  $\alpha, \beta, \gamma, \delta \in \Gamma$ .

**Proof.** Part 1. Suppose that  $\succeq^*$  is not complete so that, for some  $\alpha, \beta, \gamma, \delta \in \Gamma$ ,  $\text{Not}[(\alpha, \beta) \succeq^*(\gamma, \delta)]$  and  $\text{Not}[(\gamma, \delta) \succeq^*(\alpha, \beta)]$ . By definition, this implies  $\alpha_i a \succeq \beta_i b$ ,  $\text{Not}[\gamma_i a \succeq \delta_i b]$ ,  $\gamma_j c \succeq \delta_j d$  and  $\text{Not}[\alpha_j c \succeq \beta_j d]$ , for some  $i, j \in N$  and some  $a, b, c, d \in \mathcal{A}$ . Since  $\succeq$  is complete, we have  $\delta_i b \succeq \gamma_i a$ . Using CCI,  $\alpha_i a \succeq \beta_i b$ ,  $\delta_i b \succeq \gamma_i a$  and  $\gamma_j c \succeq \delta_j d$  imply  $\alpha_j c \succeq \beta_j d$ , a contradiction.

Part 2. Suppose that, for some  $\alpha, \beta, \gamma, \delta \in \Gamma$ ,  $\text{Not}[(\alpha, \beta) \succeq^*(\gamma, \delta)]$  and  $\text{Not}[(\beta, \alpha) \succeq^*(\delta, \gamma)]$ . By definition, we have  $\gamma_i a \succeq \delta_i b$ ,  $\text{Not}[\alpha_i a \succeq \beta_i b]$ ,  $\delta_j c \succeq \gamma_j d$  and  $\text{Not}[\beta_j c \succeq \alpha_j d]$ , for some  $i, j \in N$  and some  $a, b, c, d \in \mathcal{A}$ . Since  $\succeq$  is complete, we have  $\beta_i b \succeq \alpha_i a$ . Using CCI,  $\beta_i b \succeq \alpha_i a$ ,  $\gamma_i a \succeq \delta_i b$  and  $\delta_j c \succeq \gamma_j d$  imply  $\beta_j c \succeq \alpha_j d$ , a contradiction. Part 3 easily follows from parts 1 and 2.

Part 4. Since  $\succeq^{**}$  is complete and reversible, we have  $(\alpha, \alpha) \sim^*(\beta, \beta)$ , for all  $\alpha, \beta \in \Gamma$ . CI therefore follows using Proposition 5.

Part 5. Let  $\alpha = a_i$  and  $\beta = b_i$ . Suppose that  $a \sim b$  and  $(\gamma, \delta) \succeq^{**}(\alpha, \beta)$ . Since we have  $(\gamma, \delta) \succeq^*(\alpha, \beta)$ , we obtain, using Eq. (1),  $\gamma_i a \succeq \delta_i b$ . Suppose therefore, in contradiction with the thesis, that  $\delta_i b \succeq \gamma_i a$ . Since  $\succeq^{**}$  is complete,  $(\gamma, \delta) \succ^{**}(\alpha, \beta) \Leftrightarrow \text{Not}[(\alpha, \beta) \succeq^{**}(\gamma, \delta)] \Leftrightarrow [(\gamma, \delta) \succ^*(\alpha, \beta) \text{ or } (\beta, \alpha) \succ^*(\delta, \gamma)]$ .

Suppose that  $(\gamma, \delta) \succ^*(\alpha, \beta)$ . This implies that there are  $c, d \in \mathcal{A}$  such that, for some  $j \in N$ ,  $\gamma_j c \succeq \delta_j d$  and  $\text{Not}[\alpha_j c \succeq \beta_j d]$ . Now  $\alpha_i a \succeq \beta_i b$ ,  $\delta_i b \succeq \gamma_i a$  and  $\gamma_j c \succeq \delta_j d$  imply, using CCI,  $\alpha_j c \succeq \beta_j d$ , a contradiction. The case  $(\beta, \alpha) \succ^*(\delta, \gamma)$  is similar.

Part 6. By hypothesis, each state  $i \in N$  is influent. Therefore, there are  $\alpha, \beta, \gamma, \delta \in \Gamma$  and  $a, b \in \mathcal{A}$  such that  $\alpha_i a \succeq \beta_i b$  and  $\text{Not}[\gamma_i a \succeq \delta_i b]$  (and, hence,  $\delta_i b \succ \gamma_i a$ , since  $\succeq$  is complete). If  $\gamma_i c \succ \delta_i c$  or  $\delta_i c \succ \gamma_i c$  for some  $c \in \mathcal{A}$ , then state  $i \in N$  is essential by construction. Suppose therefore that  $\gamma_i c \sim \delta_i c$ , for all  $c \in \mathcal{A}$ .

It is easy to show that when  $\succeq$  is complete and satisfies CCI, if any of the premises of CCI holds with  $\succ$  instead of  $\succeq$ , the conclusion of CCI must hold with  $\succ$ . Now, using CCI,  $\alpha_i a \succeq \beta_i b$ ,  $\delta_i b \succ \gamma_i a$  and  $\gamma_i c \sim \delta_i c$  imply  $\alpha_i c \succ \beta_i c$ . Hence state  $i \in N$  is essential.  $\square$

As was the case with CI, it turns out that CCI can easily be characterized using our relations comparing preference differences.

**Proposition 7.** *Let  $\succeq$  be a complete relation on  $\mathcal{A}$ . Then  $\succeq$  satisfies CCI iff  $\succeq^{**}$  is complete and  $[a \sim b \text{ and } (\gamma, \delta) \succ^{**}(a_i, b_j)] \Rightarrow \gamma_i a \succ \delta_j b$ , for all  $a, b \in \mathcal{A}$ , all  $i \in \mathbb{N}$  and all  $\gamma, \delta \in \Gamma$ .*

**Proof.** Necessity results from Lemma 6. We show sufficiency. Suppose that  $\alpha_i a \succeq \beta_j b$ ,  $\gamma_i b \succeq \delta_j a$  and  $\delta_j c \succeq \gamma_j d$ . In contradiction with CCI suppose that  $\beta_j d \succ \alpha_j c$ . By definition,  $\delta_j c \succeq \gamma_j d$  and  $\text{Not}[\alpha_j c \succeq \beta_j d]$  imply  $\text{Not}[(\alpha, \beta) \succeq^*(\delta, \gamma)]$ , so that  $(\delta, \gamma) \succ^{**}(\alpha, \beta)$ , since  $\succeq^{**}$  is complete. Now  $\alpha_i a \succeq \beta_j b$  and  $(\delta, \gamma) \succ^{**}(\alpha, \beta)$  imply  $\delta_j a \succ \gamma_j b$ , a contradiction.  $\square$

The above proposition shows that a complete binary relation  $\succeq$  on  $\mathcal{A}$  that satisfies CCI is already quite well structured: there is a reversible weak order comparing preference differences between outcomes and  $\succeq$  is strictly monotonic w.r.t. this relation. The additional strength of CCI compared to CI should be apparent considering their respective impact on  $\succeq^*$  and  $\succeq^{**}$ . On its own, CI, even when coupled with completeness, does not imply that any of our relations comparing preference differences are complete.

Our experience is that the structuring of preferences brought by CCI is even more apparent considering its consequences in terms of numerical representations to which we now turn.

### 3.2. Numerical representations

We envisage a model in which:

$$a \succeq b \Leftrightarrow F(p(a_1, b_1), p(a_2, b_2), \dots, p(a_n, b_n)) \geq 0 \tag{M}$$

where  $p$  is a real-valued function on  $\Gamma^2$  that is skew symmetric (i.e.  $p(\alpha, \beta) = -p(\beta, \alpha)$ ) and  $F$  is a real-valued function on  $p(\Gamma^2)^n$  that is increasing in all its arguments and odd (i.e. such that  $F(\mathbf{x}) = -F(-\mathbf{x})$ , abusing notation in an obvious way). As shown below, model (M) turns out to have close links with CCI.

**Proposition 8.** *Suppose that  $\Gamma$  is finite or countably infinite. A binary relation  $\succeq$  on  $\mathcal{A}$  satisfies model (M) iff it is complete and satisfies CCI.*

**Proof.** Necessity. The completeness of  $\succeq$  follows from the skew-symmetry of  $p$  and the oddness of  $F$ . Suppose now that CCI is violated so that  $\alpha_i a \succeq \beta_j b$ ,  $\gamma_i b \succeq \delta_j a$ ,  $\delta_j c \succeq \gamma_j d$  and  $\text{Not}[\alpha_j c \succeq \beta_j d]$ .

Using model (M) we have, abusing notation,

$$\begin{aligned} F(p(\alpha, \beta), \mathbf{K}_{-i}) &\geq 0, \\ F(p(\gamma, \delta), -\mathbf{K}_{-i}) &\geq 0, \\ F(p(\delta, \gamma), \mathbf{L}_{-j}) &\geq 0, \\ F(p(\alpha, \beta), \mathbf{L}_{-j}) &< 0. \end{aligned}$$

Using the oddness of  $F$ , its increasingness and the skew symmetry of  $p$ , the first and the second inequalities imply  $p(\alpha, \beta) \geq p(\delta, \gamma)$ , whereas the third and the fourth imply  $p(\delta, \gamma) > p(\alpha, \beta)$ , a contradiction.

Sufficiency. Since  $\succeq$  is complete and satisfies CCI, we know from Lemma 6 that  $\succeq^{**}$  is complete so that it is a weak order. This implies that  $\succeq^*$  is a weak order.

Therefore, since  $\Gamma$  is finite or countably infinite, there is a real-valued function  $q$  such that, for all  $\alpha, \beta, \gamma, \delta \in \Gamma$ ,

$$(\alpha, \beta) \succeq^*(\gamma, \delta) \Leftrightarrow q(\alpha, \beta) \geq q(\gamma, \delta). \quad (5)$$

Using any such function  $q$ , define  $p$  letting, for all  $\alpha, \beta \in \Gamma$ ,  $p(\alpha, \beta) = q(\alpha, \beta) - q(\beta, \alpha)$ . It is easy to show that  $p$  is skew-symmetric and represents  $\succeq^{**}$ .

Define  $F$  letting, for all  $a, b \in \mathcal{A}$ ,

$$F(p(a_1, b_1), p(a_2, b_2), \dots, p(a_n, b_n)) = \begin{cases} \exp\left(\sum_{i=1}^n p(a_i, b_i)\right) & \text{if } a \succ b, \\ 0 & \text{if } a \sim b, \\ -\exp\left(-\sum_{i=1}^n p(a_i, b_i)\right) & \text{otherwise.} \end{cases} \quad (6)$$

The well-definedness of  $F$  follows from Eq. (4). It is odd by construction since  $\succeq$  is complete. Let us show that it is increasing. Suppose that  $p(\alpha, \beta) > p(a_i, b_i)$ . If  $a \succ b$ , we obtain, using Eq. (3),  $\alpha_i a \succ \beta_i b$  and the conclusion follows from the definition of  $F$ . If  $a \sim b$ , we obtain, using Proposition 7,  $\alpha_i a \succ \beta_i b$  and the conclusion follows from the definition of  $F$ . If  $b \succ a$ , we have either  $\beta_i b \succ \alpha_i a$ ,  $\beta_i b \sim \alpha_i a$ , or  $\alpha_i a \succ \beta_i b$ . In either case, the conclusion follows from the definition of  $F$ .  $\square$

**Remark 9.** Before tackling the general case, let us note that the uniqueness properties of the functions used in model (M) are clearly quite weak. Since we do not study this model for its own sake but as a framework allowing us to understand the consequences of a number of requirements on  $\succeq$ , we do not study them here; an analysis of these properties is easily inferred from the results in [Bouyssou and Pirlot \(2002\)](#).

The extension of proposition Proposition 8 to sets of arbitrary cardinality is straightforward. Let  $\langle F, p \rangle$  be a representation of  $\succeq$  in model (M). It is clear that we must have:

$$(\alpha, \beta) \succ^{**}(\gamma, \delta) \Rightarrow p(\alpha, \beta) > p(\gamma, \delta). \quad (7)$$

Hence, when model (M) holds, the weak order  $\succeq^p$  induced on  $\Gamma^2$  by  $p$  refines  $\succeq^{**}$ . A necessary condition for model (M) to hold is  $\Gamma^2$  has a finite or countable order dense subset for  $\succeq^p$  (see [Fishburn, 1970](#), ch. 3 or [Krantz et al., 1971](#), ch. 2). Since this weak order refines  $\succeq^{**}$ , it is clear that  $\Gamma^2$  will then have a finite or countable order dense subset for  $\succeq^{**}$ . Let us call OD the assertion that  $\Gamma^2$  has a finite or countable order dense subset for  $\succeq^{**}$ . We have shown that OD is necessary for model (M). The proof of Proposition 8 shows that adding this condition to the completeness of  $\succeq$  and CCI is also sufficient for (M). We omit the cumbersome and apparently uninformative reformulation of OD in terms of  $\succeq$ . We have thus proved the following.

**Theorem 10.** *A binary relation  $\succeq$  on  $\mathcal{A}$  satisfies model (M) iff it is complete and satisfies CCI and OD.*

Using model (M), it is easy to derive more consequences of the combination of completeness and CCI. We omit the simple proof of the following, which shows that, in spite of the absence of transitivity of  $\succeq$ , model (M) implies many of the usual monotonicity properties of the SEU model.

**Proposition 11.** *Let  $\succeq$  be a binary relation on  $\mathcal{A}$  satisfying model (M). Then for all  $a, b \in \mathcal{A}$ , all  $\alpha, \beta \in \Gamma$  and all nonempty  $E \subset N$ ,*

- (1)  $[a_i \succeq_{\Gamma} b_i \text{ for all } i \in N] \Rightarrow [a \succeq b]$ ,
- (2)  $[a_i \succeq_{\Gamma} b_i \text{ for all } i \in N \text{ and } a_j \succ_{\Gamma} b_j \text{ for some } j \in N] \Rightarrow [a \succ b]$ ,
- (3)  $\alpha_E a \succeq \beta_E a \Leftrightarrow \alpha \succeq_{\Gamma} \beta$ .

**Remark 12.** It may be instructive to analyze which of the classical postulates used in Savage (1954) (excluding P6 and P7, which are not pertinent in our setting with a finite number of states) are satisfied by model (M). It is not difficult to see that model (M) keeps all of P2 (since CI holds), P3 (in view of part 3 of proposition 11) and P5 (it is easy to see that  $\alpha \succ_{\Gamma} \beta \Leftrightarrow p(\alpha, \beta) > 0$ , which must be true for some  $\alpha, \beta$  if all states are to be influent; this implies that  $\succ$  cannot be empty). It keeps only the completeness part of P1, abandoning transitivity. Simple examples show that model (M) may violate postulate P4. As should be apparent from its formulation, model (M) does not, in general, allow tastes to be separated from beliefs.

**Remark 13.** Consider now model (M'), which is obtained from model (M) abandoning the increasingness of  $F$  in its arguments. In order to better appreciate the relative strengths of CI and CCI, it is interesting to note that the conjunction of completeness and CI is tantamount to (M'). We show this below in the case of a finite or countably infinite set  $\Gamma$ , leaving to the reader the, easy, task of extending the result to the general case (this requires limiting the cardinality of  $\Gamma^2 / \sim^{**}$ ).

**Proposition 14.** *Suppose that  $\Gamma$  is finite or countably infinite. A binary relation  $\succeq$  on  $\mathcal{A}$  satisfies model (M') iff it is complete and satisfies CI.*

**Proof.** Necessity. The completeness of  $\succeq$  follows from the skew-symmetry of  $p$  and the oddness of  $F$ . CI follows from  $p(\alpha, \alpha) = 0$  and  $F(\mathbf{0}) = 0$ .

Sufficiency. Let  $\geq_{\Gamma}$  be any linear order on  $\Gamma$ , i.e. a complete, antisymmetric and transitive relation. Consider the set  $\mathcal{Y} = \{(\alpha, \beta) : \alpha, \beta \in \Gamma \text{ and } \alpha >_{\Gamma} \beta\}$ , where  $>_{\Gamma}$  denotes the asymmetric part of  $\geq_{\Gamma}$ . Since  $\Gamma$  is finite or countably infinite, so is  $\mathcal{Y}$ . Therefore, there is a one-to-one function  $q$  between  $\mathcal{Y}$  and some subset of  $\mathbb{N} \setminus \{0\}$ .

Define  $p$  on  $\Gamma^2$  letting, for all  $\alpha, \beta \in \Gamma$ ,

$$\begin{aligned} p(\alpha, \beta) &= q(\alpha, \beta) \text{ if } (\alpha, \beta) \in \mathcal{Y}, \\ p(\alpha, \beta) &= 0 \text{ if } \alpha = \beta, \\ p(\alpha, \beta) &= -q(\beta, \alpha) \text{ if } \alpha \neq \beta \text{ and } (\alpha, \beta) \notin \mathcal{Y}. \end{aligned}$$

By construction,  $p$  is skew symmetric. Using such a  $p$  define  $F$  letting:

$$F(p(a_1, b_1), \dots, p(a_n, b_n)) = \begin{cases} +1 & \text{if } a \succ b, \\ 0 & \text{if } a \sim b, \\ -1 & \text{otherwise.} \end{cases} \quad (8)$$

By construction,  $p(\alpha, \beta) = p(\gamma, \delta)$  implies  $[\alpha = \beta \text{ and } \gamma = \delta]$ . Since,  $\succeq$  satisfies CI, we know from Proposition 5 that  $(\alpha, \alpha) \sim^*(\gamma, \gamma)$  so that  $(\alpha, \alpha) \sim^{**}(\gamma, \gamma)$ . This shows, using Eq. (4), that  $F$  is well-defined. It is odd since  $\succeq$  is complete.  $\square$

**Remark 15.** In Bouyssou and Pirlot (2002), we consider several models that, when translated into the framework of decision under uncertainty, fall “in between” and (M') and (M), e.g. a model in which  $F$  is odd and nondecreasing in all its arguments. The analysis of such models is straightforward adequately reformulating the conditions introduced by Bouyssou and Pirlot (2002).

#### 4. Discussion

It may be interesting to briefly compare Theorem 10 with the characterization of SEU proposed by Wakker (1989a). We recall his result below considering only the case in which there are at least two states and all states are influent (and, hence, essential—therefore we omit the condition of separability, see Wakker (1989a, Remark A.3.1, page 163) or Wakker (1988a, Th. 6.2, page 430)).

**Theorem 16.** (Wakker (1989a, Th. IV.2.7, page 83)). *Suppose that  $\Gamma$  is a connected topological space and endow  $\mathcal{A}$  with the product topology. Suppose that  $n \geq 2$  and that all states are influent. There is a continuous real valued function  $u$  on  $\Gamma$  and  $n$  positive real numbers  $p_i$  that add up to 1 such that, for all  $a, b \in \mathcal{A}$ ,*

$$a \succeq b \Leftrightarrow \sum_{i=1}^n p_i u(a_i) \geq \sum_{i=1}^n p_i u(b_i), \quad (\text{SEU})$$

*iff*

- $\succeq$  is complete,
- $\succeq$  satisfies CCI,
- $\succeq$  is continuous (i.e. the sets  $\{a \in \mathcal{A}: a \succ b\}$  and  $\{a \in \mathcal{A}: b \succ a\}$  are open for all  $b \in \mathcal{A}$ ),
- $\succeq$  is transitive.

Furthermore,  $u$  is unique up to a scale and location and the  $p_i$  are unique.

Theorem 10 abandons the topological assumptions on  $\Gamma$  and hence continuity. It also drops transitivity. Given such differences, it is rather surprising that, as already observed,

the resulting model (M) keeps a number of important properties of model (SEU). This is an indication of the power of CCI combined with completeness.

It should be noted that model (M) is far from being the only possible model taking intransitivities into account in decision making under uncertainty. Models of this type have already been suggested by Fishburn (1984, 1988, 1989, 1990), Fishburn and Lavalley (1987a,b), Lavalley and Fishburn (1987), Nakamura (1998). Most of these models are closely related to model (M) but use an additive  $F$  together with probabilities for each state. The closest to model (M) is the Skew Symmetric Additive (SSA) model with a finite number of states introduced by Fishburn (1990). This model uses the following numerical representation:

$$a \succsim b \Leftrightarrow \sum_{i=1}^n p_i \Phi(a_i, b_i) \geq 0, \quad (\text{SSA})$$

where  $p_i$  are positive real numbers that add up to one and  $\Phi$  is a skew symmetric real-valued function on  $\Gamma^2$ .

It is not difficult to see that this model implies the completeness of  $\succsim$  as well as CCI. The characterization proposed by Fishburn (1990) requires a rich topological structure for  $\Gamma$  and a formulation of continuity adapted to the nontransitive case. The necessary axioms that are used (i.e. axioms 3–5) have close connections with CCI while being collectively stronger.

## Acknowledgements

We wish to thank Peter Wakker for his very helpful comments on an earlier draft of this text. The usual caveat applies.

## References

- Abdellaoui, M., 2000. Parameter-free elicitation of utility and probability weighting functions. *Management Science* 46, 1497–1512.
- Aleskerov, F., Monjardet, B., 2002. *Utility Maximization, Choice and Preference*. Springer-Verlag, Heidelberg.
- Bleichrodt, H., Pinto, J.L., 2000. A parameter-free elicitation of the probability weighting function in medical decision making. *Management Science* 46, 1485–1496.
- Bouyssou, D., Pirlot, M., 2002. Nontransitive decomposable conjoint measurement. *Journal of Mathematical Psychology* 46, 677–703.
- Fishburn, P.C., 1970. *Utility theory for decision-making*. Wiley, New York.
- Fishburn, P.C., 1984. SSB utility theory and decision-making under uncertainty. *Mathematical Social Sciences* 8 (3), 253–285.
- Fishburn, P.C., 1985. *Interval Orders and Intervals Graphs*. Wiley, New York.
- Fishburn, P.C., 1988. *Nonlinear Preference and Utility Theory*. Johns Hopkins Univ. Press, Baltimore.
- Fishburn, P.C., 1989. Non-transitive measurable utility for decision under uncertainty. *Journal of Mathematical Economics* 18, 187–207.
- Fishburn, P.C., 1990. Skew symmetric additive utility with finite states. *Mathematical Social Sciences* 19, 103–115.
- Fishburn, P.C., 1991. Nontransitive preferences in decision theory. *Journal of Risk and Uncertainty* 4, 113–134.

- Fishburn, P.C., Lavalley, I.H., 1987a. A nonlinear, nontransitive and additive-probability model for decisions under uncertainty. *Annals of Statistics* 15 (2), 830–844.
- Fishburn, P.C., Lavalley, I.H., 1987b. State-dependent SSB utility. *Economics Letters* 25 (1), 21–25.
- Furkhen, G., Richter, M.K., 1991. Additive utility. *Economic Theory* 1, 83–105.
- Iveson, G., Falmagne, J.-C., 1985. Statistical issues in measurement. *Mathematical Social Sciences* 10, 131–153.
- Köbberling, V., Wakker, P.P., 2003. Preference foundations for nonexpected utility: a generalized and simplified technique. *Mathematics of Operations Research* 28, 395–423.
- Krantz, D.H., Luce, R.D., Suppes, P., Tversky, A., 1971. *Foundations of measurement. Additive and Polynomial Representations*, vol. 1. Academic Press, New York.
- Lavalley, I.H., Fishburn, P.C., 1987. Decision analysis under states-additive SSB preferences. *Operations Research* 35 (5), 722–735.
- Luce, R.D., 1956. Semioorders and a theory of utility discrimination. *Econometrica* 24, 178–191.
- Luce, R.D., 2000. *Utility of Gains and Losses: Measurement-Theoretical and Experimental Approaches*. Lawrence Erlbaum, Mahwah, New Jersey.
- Nakamura, Y., 1998. Skew-symmetric additive representations of preferences. *Journal of Mathematical Economics* 30, 367–387.
- Pfanzagl, J., 1971. *Theory of Measurement*, 2nd edition. Physica-Verlag, Würzburg.
- Pirlot, M., Vincke, Ph., 1997. *Semioorders. Properties, Representations, Applications*. Kluwer Academic Publishing, Dordrecht.
- Raiffa, H., 1968. *Decision Analysis—Introductory Lectures on Choices Under Uncertainty*. Addison-Wesley, Reading, Mass.
- Savage, L.J., 1954. *The Foundations of Statistics*. Wiley, New York.
- Tversky, A., 1969. Intransitivity of preferences. *Psychological Review* 76, 31–48.
- Wakker, P.P., 1984. Cardinal coordinate independence for expected utility. *Journal of Mathematical Psychology* 28, 110–117.
- Wakker, P.P., 1988a. The algebraic versus the topological approach to additive representations. *Journal of Mathematical Psychology* 32, 421–435.
- Wakker, P.P., 1988b. Derived strengths of preference relations on coordinates. *Economic Letters* 28, 301–306.
- Wakker, P.P., 1989a. *Additive Representations of Preferences: A New Foundation of Decision Analysis*. Kluwer Academic Publishing, Dordrecht.
- Wakker, P.P., 1989b. Continuous subjective expected utility with nonadditive probabilities. *Journal of Mathematical Economics* 18 (1), 1–27.
- Wakker, P.P., 1991. Additive representations of preferences, a new foundation of decision analysis; the algebraic approach. In: Doignon, J.-P., Falmagne, J.-C. (Eds.), *Mathematical Psychology: Current Developments*. Springer-Verlag, Berlin, pp. 71–87.
- Wakker, P.P., 1994. Separating marginal utility and probabilistic risk aversion. *Theory and Decision* 36 (1), 1–44.
- Wakker, P.P., Deneffe, D., 1996. Eliciting von Neumann–Morgenstern utilities when probabilities are distorted or unknown. *Management Science* 42, 1131–1150.
- Wakker, P.P., Tversky, A., 1993. An axiomatization of cumulative prospect theory. *Journal of Risk and Uncertainty* 7, 147–176.
- Wakker, P.P., Zank, H., 1999. A unified derivation of classical subjective expected utility models through cardinal utility. *Journal of Mathematical Economics* 32, 1–19.